

CAUSES AND PREVENTION OF EPILEPSY.

Amonova Zakhro Kakhramonovna, PhD., assistant of the Department of Neurology, Samarkand State Medical University
Niyozov Shukhrat Toshtemirovich Doctor of medical sciences , associate professor of the Department of Neurology of Samarkand State Medical University
Amonova Zilola Kakhramonovna is a 1st year student of the Faculty of Stomatology of the European Medical University

Abstract: Epilepsy is a neurological disorder characterized by recurrent seizures, which are sudden, uncontrolled electrical disturbances in the brain. These seizures can vary in intensity and duration, affecting a person's mind, actions, and emotions. Understanding the causes and taking preventative measures are essential to effectively managing epilepsy.

Key words: epilepsy, infections, meningitis, brain, encephalitis, medical care.

Some forms of epilepsy have a genetic component, and certain genes can predispose people to the disease. A family history of epilepsy can increase the risk of developing the disease. Head injuries, strokes, brain tumors, and infections such as meningitis can damage the brain and trigger epilepsy. Any injury that disrupts the normal electrical activity of the brain can cause seizures. Structural abnormalities in the brain, such as disorders, injuries, or developmental disorders, can cause epilepsy. These abnormalities can disrupt the normal functioning of brain cells and networks. Certain infections, such as neurocysticercosis, encephalitis, and AIDS-related infections, can cause inflammation in the brain and increase the risk of epilepsy. Conditions such as autism, Down syndrome, and neurofibromatosis are associated with an increased risk of epilepsy due to underlying brain abnormalities.[5]

Missing doses or suddenly stopping medication can trigger seizures. Maintaining a healthy lifestyle through regular exercise, a balanced diet, adequate sleep, and stress management can help reduce the risk of seizures. It is important to avoid triggers such as alcohol, drugs and sleep deprivation. Creating a safe environment by eliminating potential hazards, using safety equipment (such as helmets for high-risk activities), and educating family members and caregivers on seizure first aid can prevent injuries during a seizure. Regular visits to health care providers for follow-up and follow-up, as well as diagnostic tests such as EEG and MRI, can help effectively manage epilepsy and detect any changes in the condition. Seizure episodes, triggers, medications recording medications and lifestyle factors can provide valuable insights for health care providers to optimize treatment and identify ways to help prevent seizures can improve quality. Cooperation with health care providers, adherence to

treatment plans, and adopting a healthy lifestyle are key components in preventing and managing epilepsy. Brain injury can lead to the development of epilepsy through a variety of mechanisms that disrupt the brain's normal electrical activity and function.[1]

A brain injury, such as head trauma, stroke, tumors, infections, or other causes, can disrupt the complex network of neurons in the brain. This disorder can cause abnormal electrical impulses and neuronal over-excitability, which increases the likelihood of seizures. After brain injuries or surgery, scar tissue can form in damaged areas of the brain. This scar tissue, also called gliosis, can interfere with the transmission of electrical signals between neurons, causing abnormal electrical activity that can trigger seizures. Structural abnormalities in the brain, such as malformations, injuries, or developmental disorders that result from brain damage, can create areas of the brain that are prone to seizures. These abnormalities can change the way neurons communicate and increase the risk of epileptic activity. Brain damage can cause neurons to become overexcited and release excessive amounts of neurotransmitters, particularly glutamate, which can cause excitotoxicity. This excitotoxicity can damage neurons and disrupt the balance of neurotransmitters, contributing to the development of epilepsy. Inflammatory processes caused by brain injuries or infections can damage brain cells and tissues. Chronic inflammation and neurodegeneration can alter the normal functioning of neurons and neural circuits, creating a favorable environment for epileptic activity. Brain damage affects the function of ion channels in neurons, which are responsible for regulating the flow of ions needed to generate electrical impulses can affect.[2]

Changes in ion channel function can lead to abnormal neuronal excitability and synchronization, which increases the risk of seizures. In general, brain damage can lead to a state of neuronal hyperexcitability, where neurons behave abnormally and is more prone to firing synchronously. This hyperexcitability can manifest as an epileptic seizure when abnormal electrical activity spreads throughout the brain. Several lifestyle factors can play an important role in reducing the risk of seizures in people with epilepsy. It is important that sufferers take antiepileptic drugs as prescribed by their healthcare provider. Taking medication regularly can help control seizures and reduce the risk of episodes. Eating a balanced, nutritious diet supports overall brain health and helps manage epilepsy. Some people with epilepsy find that special diets, such as the ketogenic diet or the low-glycemic index diet, can be helpful in reducing the frequency of seizures. Sufficient and quality sleep is very important for people with epilepsy. Lack of sleep or irregular sleep patterns can trigger seizures in some people. Maintaining a regular sleep schedule and good sleep hygiene can help reduce the risk of seizures. Stress and anxiety can be triggers for seizures in some people with epilepsy. Engaging in stress-reducing activities such as mindfulness, meditation, yoga, deep

breathing exercises, or hobbies can help manage stress levels and improve overall well-being. Physical activity can have positive effects on both physical and mental health, including for people with epilepsy. Regular exercise can help reduce stress, improve mood, and improve overall fitness, which can help better control seizures. Identifying and avoiding potential seizure triggers can help prevent seizures. Common triggers include flashing lights, certain foods or drinks, lack of sleep, stress, and hormonal changes. Understanding personal triggers and taking steps to avoid them can be helpful. Alcohol and certain recreational drugs can lower the seizure threshold and increase the risk of seizures in people with epilepsy. Limiting or avoiding alcohol and illegal substances can help manage seizures.[4]

Regular visits to an epilepsy health care provider are important to monitor seizure control, adjust medications as needed, and address any concerns or changes in symptoms taking safety precautions such as avoiding activities, using protective equipment when necessary, and informing family members, friends, and colleagues about epilepsy and how to respond can help provide a seizure-safe environment. By incorporating these lifestyle factors into their daily lives, people with epilepsy can better manage their condition, reduce their risk of seizures, and improve their quality of life. For individuals with epilepsy, it is important to work closely with their health care team to develop a comprehensive treatment plan that addresses the medical and lifestyle aspects of epilepsy management.[3]

Conclusion: In conclusion, brain injury disrupts neuronal networks, creates scar tissue, alters brain structure, induces excitatory toxicity, promotes inflammation, affects ion channels, and induces neuronal hyperactivity can predispose people to epilepsy by increasing excitability. Understanding these mechanisms may help in the management and treatment of epilepsy in people with a history of brain injury.

References:

1. Ахмедов, Ш. К., Орипов, Р. А., Самандаров, Н. В., & Мамадиева, Н. Ф. ВНИМАТЕЛЬНОЕ ОТНОШЕНИЕ К АНАМНЕЗУ ПРИ ЗАБОЛЕВАНИИ ХРОНИЧЕСКОЙ КРАПИВНИЦЕЙ—ОДИН ИЗ ВАЖНЫХ ФАКТОРОВ ПРОДУКТИВНОЙ ТЕРАПИИ.[1]
2. Ахмедов, Ш. К., Абдиев, З. М., & Абдамитов, О. Р. (2013). Роль дисбактериоза кишечника при развитии атопического дерматита у детей. Академический журнал Западной Сибири,9(3),12-12.[2]
3. Ахмедов, Ш. К., Эшбаев, Э. Х., & Орипов, Р. А. (2012). ОСОБЕННОСТИ ИММУНОТЕРАПИИ ПРИ ХРОНИЧЕСКОМ ТРИХОМОНИАЗЕ. Академический журнал Западной Сибири,(4),24-25.[3]
4. Ахмедов, Ш. К., Купадзе, Р. В., Камалов, И. Ш., Абдамитов, О. Р., & Рисмурадов, Ж. М. (2014). Новое направление в клиникоиммунологическом

статусе больных ранними формами сифилиса. Академический журнал
Западной Сибири, 10(3), 78-80.[4]

5. www.ziyonet.uz [5]

